A study of electron recombination using highly ionizing particles in ArgoNeuT

Bruce Baller
Fermilab



ArgoNeut Collaboration

F. Cavanna
University of L'Aquila

A. Eriditato, S. Haug, B. Rossi, M. Weber
University of Bern

B. Baller, C. James, S. Pordes, G. Rameika, B. Rebel
Fermi National Accelerator Laboratory

M. Antonello, O. Palamara
Gran Sasso National Laboratory

T. Bolton, S. Farooq, G. Horton-Smith, D. McKee
Kansas State University

C. Bromberg, D. Edmunds, P. Laurens, B. Page
Michigan State University

J. Asaadi, Mitch Soderberg
Syracuse University

K. Lang, R. Mehdiyev
University of Texas – Austin

C. Adams, C. Anderson, E. Church, B. Fleming, R. Guenette, S. Linden, K. Partyka, J. Spitz, A. Szelc
Yale University

Contributing Author

M. Wojcik
Lodz University of Technology

Outline

- Brief survey of recombination theory
- LAr – ideal liquid vs real liquid
- Application of Birks and Box model equations
  - Introduce a modification to the Box model
- Recombination simulation
  - Focus on angular dependence
- ArgoNeuT LAr TPC in the NuMI neutrino beam
  - track and calorimetric reconstruction
- A novel(?) stopping particle ID scheme for selecting protons and deuterons
- Angular dependence – protons
- Extend to higher stopping power - deuterons

Recombination





2. *Zur Theorie der Ionisation in Kolonnen;* *von George Jaffé.*

Leipzig, im Mai 1913.

(Eingegangen 30. Mai 1913.)

$$(32') \quad Y_3(X) = \frac{1}{1 + \frac{\alpha N_0}{8\pi D}\sqrt{\frac{\pi}{z'}}\, S(z')}, \qquad z' = \frac{b^2 u^2 X^2 \sin^2\varphi}{2 D^2}.$$

Birks model (1951)

$Y_3(X)$ = recombination factor $\mathcal{R}$
→ fraction of electrons that escape vs E field strength X

Assumptions

Recombination ~ charge density

No Coulomb interactions

Ion mobility = electron mobility

Electrons & ions have the same Gaussian distribution



Liquid Argon
A Special Medium

- High electron mobility
- Electron MFP = 20 nm
- Onsager radius = 130 nm ($E_{Coulomb} = E_{thermal}$)
- No vibration levels available → ~1 nsec thermalization time
- Electrons in Coulomb field or strong external field, $\mathscr{E}$, are not in thermal equilibrium → diffusion equations not fully applicable

Box Model

Ignore electron diffusion and ion mobility in LAr

$$\mathscr{R}_{\mathrm{Box}} = \frac{1}{\xi} ln(\alpha + \xi)$$

$$\xi = k_{\mathrm{Box}} N_o / 4a^2 \mu \mathscr{E}$$

$\mathscr{E}$ = E field

Thomas & Imel, Phys Rev A 36 (1987) 614

α = 1 in the canonical model.
We allow it to vary in the recombination fits

We set $\xi = \beta$ (dE/dx) and fit β in the recombination fits

Liquid Argon
As a Real Detector Medium

Measurement — Birks form

$$\mathcal{R}_{\mathrm{ICARUS}} = \frac{A_B}{1 + k_B \cdot (dE/dx)/\mathcal{E}}$$

$A_B = 0.800 \pm 0.003$ $\neq 1$

$k_B = 0.0486 \pm 0.0006$ kV/cm (g/cm^2/MeV)

Amoruso, et al NIM A 523 (2004) 275

Impurities

Ions can attach to water molecules, screening the Coulomb field.
Debye length λ_D= distance at which screened potential E = $E_{thermal}$
λ_D= 400 – 600 nm in ArgoNeut data
Not negligible?

Measurement

Theory $\mathcal{R} \rightarrow 0$ as $\mathcal{E} \rightarrow 0$
Heavy ions: $\mathcal{R}$= 0.003
Electrons: $\mathcal{R}$= 0.35

Doke, et al Chem. Phys. Lett 115 (1985) 3434

δ-rays



Application of Birks and Box forms to reconstruction

Inverse Birks equation is < 0 at large dQ/dx

$$dE/dx = \frac{dQ/dx}{A_B/W_{ion} - k_B \cdot (dQ/dx)/\mathcal{E}}$$

Inverse Box equation is well behaved

$$dE/dx = (exp(\beta W_{ion} \cdot (dQ/dx)) - \alpha)/\beta$$

But Box model fails to match data at low dE/dx

Solution: Let $\alpha < 1$ "Modified Box Model" ala ICARUS $A_B = 0.8$

Example with $\alpha = 0.93$ $\beta = 0.32$



Recombination Simulation

E field



Stopping proton: dE/dx = 24 MeV/cm → r_k = 10 nm

MIP: dE/dx = 1.7 MeV/cm → r_k = 50 nm

Initial conditions:
r_o = 0.5 nm, E_{ko} = 5 eV

After thermalization:
$\langle r_o \rangle$ ~ 2500 nm, $\langle E_{ko} \rangle$ ~ 0.01 eV

Simulation includes motion due to (periodic) Coulomb field, external E field and atomic collisions, escape and recombination criteria



Jaskolski, Wojcik J. Phys. Chem. A 115 (2011) 4317

Recombination Simulation

Angular Dependence

Modify simulation to allow non-perpendicular $\mathcal{E}$ field

Simulation runs for r_k = 10, 20, 30, 40, 50 nm and $\phi = 40^o, 50^o, 60^o, 80^o$

Ratios of escape probability, $\mathcal{R}$, vs dE/dx →

Simulation (data points) $\mathcal{R}_{ICARUS}$ with $\mathcal{E} \rightarrow \mathcal{E} \sin\phi$ (curves)

<u>Significant angular dependence expected from theory and simulation</u>



M. Wojcik

ArgoNeuT





Cryostat Volume	500 Liters
TPC Volume	175 Liters
# Electronic Channels	480
Electronics Style (Temp.)	JFET (293 K)
Wire Pitch (Plane Separation)	4 mm (4 mm)
Electric Field	481 V/cm
Max. Drift Length (Time)	0.5 m (330 μs)
Wire Properties	0.15mm diameter BeCu

C. Anderson, 2012 JINST 7 P10019

Stopping Particle Stopping Power

Bethe-Bloch eqn has power law dependence with residual range (R) near the stopping point

$$(dE/dx)_{hyp} = A\,R^{b}$$

$$T_{range} = \frac{A}{b+1}\,R^{b+1}$$

Particle	A	b
pion	7.9	-0.37
kaon	13.5	-0.41
proton	17	-0.42
deuteron	25	-0.43

T_{range} (MeV), R (cm)



Note the weak dependence on b

Select Highly Ionizing Particles



1) Reconstruct 3D tracks = cluster of 3D space points each with a measurement of charge Q deposited using the area of a Gaussian fit (collection plane)
2) Find dQ/dx using angle corrected distance between space points
3) Correct for electron lifetime
4) Find $(dE/dx)_{calo}$ using Birks or Box equation
5) Sum up to find kinetic energy deposited = T_{calo}
6) Find T_{range} using track length assuming a proton hypothesis
7) Eliminate lightly ionizing ptcls by requiring $T_{calo} > 0.5\ T_{range}$

Particle Identification

with minimal bias

Algorithm

Set b = constant = 0.42

Find $A_i = (dE/dx)_{calo} \times R^{0.42}$ for each space point i on a track

Define PIDA = $< A_i >$ = average value for the track

Histogram PIDA and look for bumps →

Requirements

Protons: 14 < PIDA < 21

Deuterons: 25 < PIDA < 33



2900 proton candidates

170 deuteron candidates






Range
Entries 2903
Mean 9.928
RMS 6.455
ArgoNeut data
Proton candidates
50 MeV < T_{range} < 250 MeV
Events
Range (cm)


Events
40°
50°
60°
80°
ϕ (degrees)

Angular Dependence Protons



No recombination model assumptions required to make this plot

$(dE/dx)_{hyp}$ from Bethe-Bloch using R

Angular Dependence Protons



Significantly weaker than expected from theory and simulation

Debye-like screening effect?

Recombination Fits



Vertical bars include 2% systematic error
Horizontal bars δR = 1 mm



Data – open circles
Birks fit – red curve
Modified Box fit – blue curve

Fit Summary Protons

Excellent agreement with ICARUS in the 80° bin

Angle Bin	Angle Bin Range	Box α	Box β $(MeV/cm)^{-1}$	Birks A_{Argo}	Birks k_{Argo} $(kV/cm)(g/cm^2)/MeV$
80°	70° - 90°	0.91 ± 0.03	0.302 ± 0.005	0.793 ± 0.018	0.049 ± 0.002
60°	55° - 70°	0.92 ± 0.04	0.317 ± 0.006	0.794 ± 0.019	0.052 ± 0.003
50°	47° - 55°	0.90 ± 0.04	0.327 ± 0.007	0.791 ± 0.020	0.053 ± 0.003
40°	20° - 47°	1.06 ± 0.05	0.346 ± 0.007	0.875 ± 0.025	0.066 ± 0.004

α ~ independent of angle

$\langle\alpha\rangle = 0.93 \pm 0.02$



Deuterons



Deuteron: $(dE/dx)_{hyp} = 25\ R^{-0.43}$

Summary

- Introduced a modified Box model
  - Excellent agreement with data and Birks model
  - Obviates the poor behavior of the Birks model at low ionization
- Significant recombination angular dependence expected from columnar theory and simulation
  - ~25% loss of charge collected at $\phi \sim 40^{\circ}$ and dE/dx ~ 24 MeV/cm compared to the same track at $\phi \sim 80^{\circ}$
- Introduced a PID scheme using the power-law behavior of stopping particle stopping power
- Charge loss is 5% - 10% in the proton sample at high dE/dx and small angle
- Extend the range of validity to 35 MeV/cm using a small sample of deuterons

Backup Slides



Lower PIDA range: 23 < PIDA < 27
Contaminated by protons in the Gaussian tail
See slide 14

Deuteron: $(dE/dx)_{hyp} = 25\ R^{-0.43}$



Are the deuteron candidates really protons?

Use the (incorrect) proton hypothesis with the deuteron sample

Estimating the Stopping Point Position

- The stopping point is usually assumed to be dx/2, where dx is the distance between the last two space points
- Use the pattern of dE/dx in the last 5 space points to estimate the stopping point in the last wire cell
- Step a distance δR in the last cell ($1 < \delta R < 10$ mm) in 1 mm increments
- For each step, calculate
  - $dQ/\delta R$ for the stopping point,
  - Recombination correction → $(dE/\delta R)_{calo}$ for the stopping point,
  - $(dE/dx)_{hyp}$ for next 4 points using residual range = $\delta R + n\ dx$ (n = 1,2,3,4)
- Find rms difference between $(dE/dx)_{hyp}$ and $(dE/dx)_{calo}$ for all points
- Use the Δ value with the smallest rms

Estimating the Stopping Point Position

Assume the track stops halfway in the last cell



<dE/dx> ~ 45 ± 45 MeV/cm

After fitting to the stopping point



<dE/dx> ~ 25 ± 13 MeV/cm

Estimating the Stopping Point Position

- Stopping point error from Monte Carlo ~ 1 mm
- $(dE/dx)_{calo}$ from the last point is not included in the recombination fits
- Stopping point fit reduces the $(dE/dx)_{hyp}$ error propagated from the equation on slide 12
  - Horizontal error bars on slide 18